


The Importance of Considering Mineralocorticoid Receptor (MR) Overactivation in Patients With Chronic Kidney Disease (CKD) Associated With Type 2 Diabetes (T2D)

In adults with CKD associated with T2D, there are **3 main factors** that may contribute to CKD progression¹⁻³

HEMODYNAMIC^{1,2}
Elevated BP




INFLAMMATION AND FIBROSIS IN THE KIDNEYS¹⁻⁴

Inflammation and fibrosis can contribute to permanent structural damage in the kidneys

METABOLIC^{1,2}
Elevated HbA1c

BP, blood pressure; HbA1c, glycated hemoglobin.



Consider each of the 3 main factors associated with CKD progression in patients with T2D for appropriate disease management^{1-3,6}



Encourage your patients to learn more about CKD associated with T2D by visiting

www.ckdandt2d.com

References: 1. Alicic RZ et al. *Clin J Am Soc Nephrol.* 2017;12(12):2032-2045. 2. Mora-Fernández C et al. *J Physiol.* 2014;592(18):3997-4012. 3. Bauersachs J et al. *Hypertens.* 2015;65(2):257-263. 4. Thomas MC et al. *Nat Rev Dis Primers.* 2015;1:15018. 5. Tesch GH, Young MJ. *Front Pharmacol.* 2017;8:313. 6. American Diabetes Association. *Diabetes Care.* 2022;45(suppl 1):S1-S264.

© 2022 Bayer. BAYER and the Bayer Cross are registered trademarks of Bayer. All rights reserved. MAC-UN-CAR-US-0080-1 02/22

